General Screenings and Immunizations

For **Females** Ages 18-49* at Average Risk for Most Diseases

Screening Tests	When	Why
General Health:		
Full checkup, including height and weight	Every 2-3 years. Discuss with your doctor or nurse.	To screen for diseases, assess risk for future problems, discuss lifestyle habits, and keep vaccinations up to date.
Thyroid Test (TSH)	Discuss with your doctor or nurse.	To identify an under- or overactive thyroid, both of which are very treatable and either of which can lead to more serious conditions if left untreated.
Heart Health:		
Blood Pressure Test	At least every 2 years.	To test for high blood pressure, which can lead to heart attack, stroke, heart failure, and kidney and eye problems.
Cholesterol Test	Every 5 years starting at age 35. If you smoke or are overweight, you may need to start at age 20. Discuss with doctor or nurse about when you should begin.	Treating cholesterol abnormalities can help reduce your risk of heart disease.

Screening Tests	When	Why
Bone Health:		
Bone Mineral Density Test	Talk with your doctor or nurse.	There are no obvious signs of weak bones until you fracture a bone. Bone density screenings identify problems early, allowing you to start treatment and prevent further bone loss.
Diabetes:		
Fasting Blood Sugar (Glucose) Test	Every 3 years starting at age 45. Talk with your doctor if you are at risk for diabetes or are overweight.	To provide an early warning sign of high blood sugar levels, which could mean an increased risk for diabetes.
Breast Health:		
Mammogram (X-ray of breast)	Every 1-2 years, starting at age 40.	To identify possible early signs of breast cancer.
Breast Exam	Females in their 20s and 30s should receive a clinical breast exam every 3 years, and every year for females 40 and over. Starting at age 20, females should perform breast self-exams once a month.	Not all breast cancers are found on mammograms; a good clinical breast exam and self-exam can also help identify cancers relatively early.

Screening Tests	When	Why
Reproductive Health:		
Pap Test	Females ages 21-30 should get a Pap test every 3 years. Females over 30 should get a Pap and HPV test every 3-5 years.	Helps identify females at risk for developing cervical cancer.
HPV Test (Human Papillomavirus)	Every 5 years for females ages 30-65. Should be done as part of the Pap test.	The HPV test in combination with the Pap test is better at identifying females at risk for cervical cancer, than the Pap test alone.
Pelvic Exam	Every year for females starting at age 21. Discuss with your doctor or nurse if you should start earlier.	A way to assess health, lifestyle and health risks.
Chlamydia Test	Yearly until age 25 if sexually active. For ages 26 and over, get the test if you have new or multiple sexual partners, or at higher risk.	Prevents spread of chlamydia.
Sexually Transmitted Illnesses (STI) Tests	All sexually active females and their partners should be tested for STIs, including HIV, before beginning sexual activity.	Prevents spread of HIV and other STIs, many of which can only be detected through testing.

Screening Tests	When	Why
Eye & Ear Health:		
Eye Exam	At least once from ages 20-29; at least twice between ages 30-39. Get a baseline eye disease screening at age 40. Discuss your eye health with a doctor.	To test your vision and screen for glaucoma and macular degeneration, two common, often age- related conditions.
Hearing Test	Every 10 years from ages 18-50.	To make sure you're hearing all life has to offer.
Oral Health:		
Dental Exam	1-2 times a year.	To remove plaque and bacteria that could lead to tooth and gum disease; to check for tongue and mouth cancer.
Skin Health:		
Skin Exam	Starting at age 18, females should do a monthly skin self-exam. Speak to your doctor or nurse about how often to have an additional skin exam by your doctor.	To track worrisome moles, identify skin cancer early, and be able to report any changes to your doctor or nurse.

Screening Tests	When	Why
Mental Health:		
Mental Health Screening	Discuss with your doctor or nurse.	If you are feeling sad or anxious for a long time, it might be helpful to speak with a doctor.
Immunizations:		
Influenza (Flu) Vaccine	Annually for everyone 6 months and older.	Protects against some flu viruses.
Human Papillomavirus Vaccine (HPV)	Age 11-12, or 13-26 if not previous vaccinated; 3 doses at 0-, 2-, and 6- month intervals; no booster necessary.	Protects against four common types of HPV, including the two most likely to cause cervical cancer.
Meningococcal Vaccine	College freshmen, military recruits, and other at-risk persons; discuss with your doctor or nurse.	Protects against some types of meningococcal disease (meningitis).
Tetanus, Diphtheria, Pertussis Booster Vaccine (Td/Tdap)	Every 10 years.	Protects against tetanus, diphtheria, and pertussis.
Varicella (Chickenpox)	Given in 2 doses at 0- and 4- to 8-week intervals to those 19 or older who have not been vaccinated or had chickenpox.	Protects against chickenpox, a usually mild but highly contagious childhood disease, which can be dangerous in infants and adults.

* Please Note: These charts are guidelines only. Your doctor or nurse will personalize the timing of each test and immunization to best meet your health care needs.