



Oregon Health & Science University
Hospital and Clinics Provider's Orders

PO7071



ADULT AMBULATORY INFUSION ORDER
**Agalsidase Beta (FABRAZYME)
Infusion**

Page 1 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Weight: _____ kg Height: _____ cm

Allergies: _____

Diagnosis Code: _____

Treatment Start Date: _____ Patient to follow up with provider on date: _____

****This plan will expire after 365 days at which time a new order will need to be placed****

GUIDELINES FOR ORDERING

1. **Send FACE SHEET and H&P or most recent chart note.**
2. Indicated for use in patients with Fabry disease. Reduces globotriaosylceramide (GL-3) deposition in capillary endothelium of the kidney and certain other cell types
3. Please encourage patients to enroll in the Fabry registry by visiting www.fabryregistry.com or calling 1-800-745-4447.
4. Patients with advanced Fabry disease may have compromised cardiac function which may predispose them to a higher risk of severe complications from infusion reactions.

NURSING ORDERS:

1. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, dec clotting (alteplase), and/or dressing changes.
2. VITAL SIGNS – Monitor and record vital signs, tolerance, and presence of infusion-related reactions prior to infusion, with every rate increase, then hourly until infusion is complete.
3. Initial infusion: Initial rate should not exceed 30 mL/hr.
Subsequent infusions: If no previous infusion reaction, rate may be increased in increments of 10 mL/hour with each subsequent infusion.
Maximum total infusion time: 1.5 hours (MAX RATE = total volume/1.5).
If previous infusion reaction contact provider for guidance.
4. Observe patient for 60 minutes following infusion (unless prescriber indicates this is not necessary).
5. Reschedule patient for next weekly infusion.

PRE-MEDICATIONS: (Administer 30 minutes prior to infusion)

Note to provider: Please select which medications below, if any, you would like the patient to receive prior to treatment by checking the appropriate box(s)

- acetaminophen (TYLENOL) tablet, 650 mg, oral, ONCE, every visit
- diphenhydrAMINE (BENADRYL) capsule, 50 mg, oral, ONCE, every visit
- loratadine (CLARITIN) tablet, 10 mg, oral, ONCE, every visit

(Choose as alternative to diphenhydrAMINE if needed)



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
**Agalsidase Beta (FABRAZYME)
Infusion**

Page 2 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

MEDICATIONS:

Agalsidase beta (FABRAZYME) 1 mg/kg = _____ mg in sodium chloride 0.9% IV infusion, ONCE, every 2 weeks x _____ doses (*Pharmacist will round dose to nearest 5 mg vial and modify during order verification*)

Administer using an in-line low protein binding 0.2 micron filter.

HYPERSENSITIVITY MEDICATIONS:

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in: Oregon _____ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is # _____ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: _____ **Date/Time:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
**Agalsidase Beta (FABRAZYME)
Infusion**

Page 3 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

OLC Central Intake Nurse:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

Please check the appropriate box for the patient's preferred clinic location:

Beaverton

OHSU Knight Cancer Institute

15700 SW Greystone Court
Beaverton, OR 97006

Phone number: 971-262-9000

Fax number: 503-346-8058

NW Portland

Legacy Good Samaritan campus

Medical Office Building 3, Suite 150
1130 NW 22nd Ave.
Portland, OR 97210

Phone number: 971-262-9600

Fax number: 503-346-8058

Gresham

Legacy Mount Hood campus

Medical Office Building 3, Suite 140
24988 SE Stark
Gresham, OR 97030

Phone number: 971-262-9500

Fax number: 503-346-8058

Tualatin

Legacy Meridian Park campus

Medical Office Building 2, Suite 140
19260 SW 65th Ave.
Tualatin, OR 97062

Phone number: 971-262-9700

Fax number: 503-346-8058

Infusion orders located at: www.ohsuknight.com/infusionorders